arXiv:0710.2386v1 [math.MG] 12 Oct 2007

Local Convexity Properties of j-metric Balls

Riku Klén

Abstract

This paper deals with local convexity properties of the j-metric. We consider convexity and starlikeness of the j-metric balls in convex, starlike and general subdomains of $\mathbb{R}^n$.

2000 Mathematics Subject Classification: Primary 30F45, Secondary, 30C65

Key words: j-metric ball, local convexity

`File: jmetric.tex, 2007-01-10, printed: 2008-2-2, 3.36`

1 Introduction

The *j-distance* in a proper subdomain G of the Euclidean space $\mathbb{R}^n$, $n \geq 2$, is defined by

$$j_G(x,y) = \log\left(1 + \frac{|x-y|}{\min\{d(x), d(y)\}}\right),$$

where $d(x)$ is the Euclidean distance between x and ∂G. If the domain G is understood from the context we use notation j instead of j_G.

The j-distance was first introduced by F.W. Gehring and B.P. Palka [GP] in 1976 in a slightly different form and in the above form, by M. Vuorinen [Vu2] in 1985. The j-distance is actually a metric and a proof of the triangle inequality valid for general metric spaces is given in [S]. Previously the j-metric has been studied in connection with the study of other metrics [GO, H, S, V, Vu2]. See also recent papers [HL, L]. In spite of these studies many basic questions of the j-metric remain open and some of them will be studied here.

The purpose of this paper is to study metric spaces (G, j_G) and especially local convexity properties of *j-metric balls* or in short *j-balls* defined by

$$B_j(x,M) = \{y \in G \colon j(x,y) < M\},$$

where $M > 0$ and $x \in G$. In the dimension $n = 2$ we call these *j-metric disks* or *j-disks.*

M. Vuorinen suggested in [Vu4] a general question about the convexity of balls of small radii in metric spaces. This paper is motivated by this question and we will provide an answer in a particular case. Our main result is the following theorem. For the definition of starlike domains see 3.3.

Theorem 1.1. *For a domain $G \subsetneq \mathbb{R}^n$ and $x \in G$ the j-balls $B_j(x, M)$ are convex if $M \in (0, \log 2]$ and strictly starlike with respect to x if $M \in \big(0, \log(1+\sqrt{2})\big]$.*

In Section 2 we consider general properties of the j-metric and show that for any G there exists points such that there is no geodesic between them. In Section 3 we consider local convexity properties of j-balls in punctured space and in Section 4 we extend these results to an arbitrary domain $G \subsetneq \mathbb{R}^n$. We will further consider convexity of j-balls in convex domains and starlikeness of j-balls in starlike domains.

2 Properties of the j-metric

Throughout this paper $G \subsetneq \mathbb{R}^n$, $n \geq 2$, is a domain. We denote $m(x, y) = \min\{d(x), d(y)\}$ and we use notation $B^n(x, M)$ for the Euclidean balls and $S^{n-1}(x, M)$ for the Euclidean spheres. We often identify $\mathbb{R}^2$ with the complex plane $\mathbb{C}$.

In 1976 F.W. Gehring and B.P. Palka [GP] also introduced the quasihyperbolic metric, which has been widely applied in geometric function theory and mathematical analysis in general, see e.g. [Vu3, V]. The *quasihyperbolic distance* between two points x and y in a proper subdomain G of the Euclidean space $\mathbb{R}^n$, $n \geq 2$, is defined by

$$k_G(x, y) = \inf_{\alpha \in \Gamma_{xy}} \int_\alpha \frac{|dx|}{d(x)},$$

where Γ_{xy} is the collection of all rectifiable curves in G joining x and y. We denote the *quasihyperbolic ball* by

$$D_G(x, M) = \{y \in G \colon k_G(x, y) < M\}.$$

The quasihyperbolic metric is closely related with the j-metric. By [GP, Lemma 2.1] j_G is always a minorant of k_G, in other words, for a proper subdomain G of $\mathbb{R}^n$ we have

$$j_G(x, y) \leq k_G(x, y)$$

for all $x, y \in G$.

The following result can be used to estimate the quasihyperbolic metric from above by the j-metric.

Proposition 2.1. *[Vu3, Lemma 3.7] Let* $G \subsetneq \mathbb{R}^n$ *be a domain,* $x \in G$, $y \in B^n\big(x, d(x)\big)$ *and* $s \in (0,1)$. *Then*

$$k_G(x,y) \le \frac{1}{1-s} j_G(x,y).$$

The following lemma gives Euclidean bounds for the j-balls.

Proposition 2.2. *[S, Theorem 3.8] For a proper subdomain* $G \subset \mathbb{R}^n$, $x \in G$ *and* $M > 0$ *we have*

$$B^n\big(x, r\, d(x)\big) \subset B_j(x, M) \subset B^n\big(x, R\, d(x)\big),$$

where $r = 1 - e^{-M}$ *and* $R = e^M - 1$. *Moreover*

$$B_j(x, M) \subset \big\{ z \in G \colon e^{-M} d(x) \le d(z) \le e^M d(x) \big\}.$$

Remark 2.3. A similar result to Proposition 2.2 is also true for the quasihyperbolic metric see [Vu1, page 347].

By Proposition 2.2 the j-ball $B_j(x, M)$ shrinks towards the center $\{x\}$ as M approaches 0. The following lemma shows that the j-balls $B_j(x, M)$ exhaust the domain G.

Lemma 2.4. *Let* $G \subset \mathbb{R}^n$ *be a bounded domain and fix* $x \in G$ *and* $s \in (0, d(x)]$. *Then*

$$\{y \in G \colon d(y) > s\} \subset B_j\big(x, \log(1 + d/s)\big),$$

for $d = \sup\limits_{z \in \partial G} |x - z|$.

Proof. Let us assume $d(y) > s$. Then either $m(x,y) = d(x) \ge s$ or $m(x,y) = d(y) > s$. In both cases $m(x,y) \ge s$ and since $|x - y| < d$ for all $y \in G$ we have

$$j(x,y) = \log\left(1 + \frac{|x-y|}{m(x,y)}\right) < \log\left(1 + \frac{d}{s}\right).$$

□

Let us denote the set of closest boundary points of a point x in a domain $G \subset \mathbb{R}^n$ by

$$R_x = \{z \in \partial G \colon |z - x| = d(x)\}\,.$$

The next result characterizes the case of equality in the triangle inequality for the j-metric. Its proof is based on the proof of the triangle inequality [S, Lemma 2.2].

Theorem 2.5. *Let $x, y, z \in G \subsetneq \mathbb{R}^n$ be distinct points and $d(x) \le d(z)$. Then*

$$j_G(x,z) = j_G(x,y) + j_G(y,z)$$

implies that x, z and u are collinear for some $u \in R_x$ and $y \in (x,z)$ with $d(x) < d(y) < d(z)$.

Proof. By definition $j_G(x,z) < j_G(x,y) + j_G(y,z)$ is equivalent to

$$\frac{|x-z|}{m(x,z)} < \frac{|x-y|}{m(x,y)} + \frac{|y-z|}{m(y,z)} + \frac{|x-y||y-z|}{m(x,y)m(y,z)}. \tag{2.6}$$

The assumption $d(x) \le d(z)$ implies $m(x,z) = d(x)$.

If $d(y) \le d(x)$, then the inequality (2.6) is equivalent to

$$|x-z| < |x-y|\frac{d(x)}{d(y)} + |y-z|\frac{d(x)}{d(y)} + \frac{|x-y||y-z|}{d(y)}\frac{d(x)}{d(y)},$$

which is true, because $|x-z| \le |x-y| + |y-z|$, $(|x-y||y-z|)/d(y) > 0$ and $d(x)/d(y) \ge 1$.

If $d(y) > d(x)$, then the inequality (2.6) is equivalent to

$$|x-z| < |x-y| + |y-z|\left(\frac{d(x) + |x-y|}{m(y,z)}\right),$$

which is false if and only if x, y and z are collinear and

$$\frac{d(x) + |x-y|}{m(y,z)} = 1.$$

If $d(x) = d(z)$, then $d(x)/m(y,z) = 1$ and

$$\frac{d(x) + |x-y|}{m(y,z)} > 1. \tag{2.7}$$

If $d(x) < d(z) < d(y)$, then the inequality (2.7) is true, because $d(x) + |x-y| \ge d(y) > d(z) = m(y,z)$. If $d(x) < d(y) \le d(z)$, then the inequality (2.7) is true if and only if $y \notin \{k(x-u)\colon k > 0\}$, where $u \in R_x$. $\square$

The implication of Theorem 2.5 in the other direction was proved by Hästö, Ibragimov and Lindén [HIL, Corollary 3.7].

Definition 2.8. Let $G \subsetneq \mathbb{R}^n$ be a domain and γ a curve in G. If

$$j(x,y) + j(y,z) = j(x,z)$$

for all $x, z \in \gamma$ and $y \in \gamma'$, where γ' is the subcurve of γ joining x and z, then γ is a *geodesic segment* or shortly a *geodesic*. We denote a geodesic between x and y by $J[x,y]$.

By Theorem 2.5 and the result of Hästö, Ibragimov and Lindén we can easily find all geodesics $J[x,y]$ for any domain G. The geodesic needs to satisfy the triangle inequality as equality at each point and therefore the geodesic can only be a line segment l with the following property.

Lemma 2.9. *Let $G \subsetneq \mathbb{R}^n$ be a domain and $J[x,y]$ be a geodesic segment with $x, y \in G$. There exists $u \in \partial G$ such that $u \in R_s$ for all $s \in J[x,y]$ and u and $J[x,y]$ are collinear.*

Proof. Let us assume, on the contrary, that there exists $z \in J[x,y]$ such that $d(z) < d(x) - |x-z|$. Now $j_G(x,z) + j_G(z,y) = j_G(x,y)$ is equivalent to

$$d(z)|x-z| + \big(d(x) + |x-z|\big)|z-y| = d(z)|x-y|.$$

We have

$$\begin{array}{rcl} d(z)|x-y| & \leq & d(z)|x-z| + d(z)|z-y| \\ & < & d(z)|x-z| + \big(d(x) + |x-z|\big)|z-y| \\ & = & d(z)|x-y| \end{array}$$

which is a contradiction. $\square$

Theorem 2.10. *Let $G \subsetneq \mathbb{R}^n$ be a domain. Then there exist $x, y \in G$ such that there is no geodesic $J[x,y]$.*

Proof. Let us assume, on the contrary, that for all $x, y \in G$ there exists a geodesic $J[x,y]$. Since G is a domain, we can choose $x, y, z \in G$ to be three distinct noncollinear points. Now there exists a geodesic $J[x,y]$ from x to y. We may assume $d(x) < d(y)$ and then by Lemma 2.9 $B^n\big(x, d(x)\big) \subset B^n\big(y, d(y)\big) \subset G$.

On the other hand, there exists a geodesic $J[x,z]$ from x to z and therefore there has to exist a point $u \in S^{n-1}\big(x, d(x)\big) \cap \partial G$ such that x, z and u are collinear. This is a contradiction, because x, y and u are noncollinear and therefore $u \in B^n\big(y, d(y)\big)$. $\square$

Remark 2.11. By Theorem 2.10 a j-metric geodesic does not always exist between two points. F.W. Gehring and B.G. Osgood have proved [GO, Lemma 1] that for the quasihyperbolic metric there always exists a geodesic between two points of a domain $G \subsetneq \mathbb{R}^n$.

However, the geodesics of the j-metric are unique while the geodesics of the quasihyperbolic metric need not be unique.

3 Convexity and starlikeness of j-balls in punctured space

In this section we consider the case $G = \mathbb{R}^n \setminus \{0\}$. By definition the j-balls in punctured space $G = \mathbb{R}^n \setminus \{0\}$ are similar, which means that $B_j(x,M)$ can be mapped onto $B_j(y,M)$ for all $x, y \in G$ by rotation and stretching. We see easily that these balls are also symmetric along the line that goes through 0 and the center point.

Theorem 3.1. *Let $x \in \mathbb{R}^n \setminus \{0\}$. Then*
1) the j-ball $B_j(x,M)$ is convex if and only if $M \in (0, \log 2]$.
2) the j-ball $B_j(x,M)$ is strictly convex if and only if $M \in (0, \log 2)$.

Proof. 1) By similarity we can assume $x = e_1$ and by symmetry it is sufficient to consider only the case $n = 2$. We will consider $\partial B_j(1,M)$ for fixed M. By definition we have for $z \in \partial B_j(1,M)$

$$M = \begin{cases} \log(1+|z-1|), & |z| \geq 1, \\ \log\left(1+|z-1|/|z|\right), & |z| < 1, \end{cases}$$

which is equivalent to

$$e^M - 1 = \begin{cases} |z-1|, & |z| \geq 1, \\ |1-1/z|, & |z| < 1. \end{cases}$$

For $|z| \geq 1$ the $\partial B_j(1,M)$ is an arc of a circle with center 1 and radius $e^M - 1$. For $|z| < 1$ the $\partial B_j(1,M)$ is a circle that goes through points $1/(e^M)$ and $1/(2-e^M)$ and has center on the real axis. This means that the center of the circle is $c = 1/\big(e^M(2-e^M)\big)$ and the radius of the circle is $|e^M - 1|/|e^M(2-e^M)|$. Now $c > 1$, if $M \leq \log 2$, and $c < 0$, if $M > \log 2$. Therefore $\partial B_j(1,M)$ is convex for $M \leq \log 2$ and not convex for $M > \log 2$.

2) We have $c \in (1, \infty)$, where c is as above. Therefore $B_j(x,M)$ is strictly convex. In the case $M = \log 2$ we have $c = \infty$ and $B_j(x,M)$ is not strictly convex. $\square$

Remark 3.2. For fixed $x \in G$ the quasihyperbolic ball $D_G(x, M)$ is strictly convex in $G = \mathbb{R}^n \setminus \{0\}$ if and only if $M \in (0, 1]$ [K].

Clearly $B_j(x, M)$ is never smooth. We will next define starlikeness of a domain.

Definition 3.3. Let $G \subset \mathbb{R}^n$ be a bounded domain and $x \in G$. We say that G is *starlike with respect to* x if each line segment from x to $y \in G$ is contained in G. The domain G is *strictly starlike with respect to* x for $x \in G$ if each ray from x meets ∂G at exactly one point.

The next theorem determines the values of M for which the j-ball $B_j(x, M)$ is strictly starlike with respect to x.

Theorem 3.4. *For $x \in \mathbb{R}^n \setminus \{0\}$ the j-ball $B_j(x, M)$ is strictly starlike with respect to x if and only if $M \in \big(0, \log(1 + \sqrt{2})\big]$.*

Proof. Because the j-balls are similar it is sufficient to consider $x = e_1$. By symmetry it is sufficient to consider the case $n = 2$ and the part of $\partial B_j(1, M)$ that is above the real axis. If $M \geq \log 3$, then $B_j(1, M) = B^2(1, r) \setminus B^2(c, s)$, where c, r and s are given in the proof of Theorem 3.1 and $B^2(c, s) \subset B^2(1, r)$. Therefore $B_j(1, M)$ can be starlike with respect to 1 only for $M < \log 3$.

Let us assume $M < \log 3$. By the proof of Theorem 3.1 $B_j(1, M) = B^2(1, r) \setminus B^2(c, s)$. Let us denote the point of intersection of $S^1(1, r)$ and $S^1(c, s)$ above the real axis by z. Now z is also the point of intersection of the unit circle and the boundary $\partial B_j(1, M)$. Let us denote by l the line that goes through points 1 and z. Now $B_j(1, M)$ is strictly starlike with respect to 1 if and only if $l \cap B^2(1, r) \cap B^2(c, s) = \emptyset$. If z is a tangent of $S^1(c, s)$, then the circles $S^1(1, r)$ and $S^1(c, s)$ are perpendicular and M has the largest value such that $B_j(1, M)$ is starlike with respect to 1.

By the proof of Theorem 3.1 we have $c = -1/e^M(e^M - 2)$, $r = |1 - z| = e^M - 1$, $|1 - c| = (e^M - 1)^2/e^M(e^M - 2)$ and $s = |z - c| = (e^M - 1)/e^M(e^M - 2)$. Let us assume that z is a tangent of $S^1(c, s)$. Now by the Pythagorean Theorem

$$\frac{(e^M - 1)^4}{e^{2M}(e^M - 2)^2} = (e^M - 1)^2 + \frac{(e^M - 1)^2}{e^{2M}(e^M - 2)^2},$$

which is equivalent to $e^{2M} - 2e^M - 1 = 0$ and therefore

$$M = \log(1 + \sqrt{2}).$$

$\square$



Figure 1: The boundaries of j-disks $j(1,M)$ in punctured plane $G=\mathbb{R}^2\setminus\{0\}$ with $M=0.5$, $M=\log 2$, $M=\log(1+\sqrt{2})$ and $M=1.1\approx\log 3$.

Example 3.5. Let us consider the starlikeness of j-balls $B_j(x,M)$ with respect to $z\in B_j(x,M)$ for $M>\log 2$. By choosing $z=(e^{-M}+\varepsilon)x/|x|$ for $\varepsilon>0$ and letting ε approach to zero we see that $B_j(x,M)$ is not starlike with respect to z.

On the other hand, if we choose $z=(e^M-\varepsilon)x/|x|$ for $\varepsilon>0$ and $M<\log\big((3+\sqrt{5}/2\big)$, we see that $B_j(x,M)$ is strictly starlike with respect to z for small enough ε.

Remark 3.6. For fixed $x\in G$ the quasihyperbolic ball $D_G(x,M)$ is strictly starlike with respect to x in $G=\mathbb{R}^n\setminus\{0\}$ if and only if $M\in(0,\kappa]$ [K], where $\kappa\approx 2.83297$.

4 Convexity and starlikeness of j-balls

We will consider convexity and starlikeness of j-balls $B_j(x,M)$ for $M>0$ in convex, starlike and general domains.

Let us consider j-balls in a domain G with a finite number of boundary points. The case card $\partial G=1$ is identical to $G=\mathbb{R}^n\setminus\{0\}$. If $\partial G=\{y_1,y_2\}$, then $B_{j_G}(x,M)=B_{j_{\mathbb{R}^n\setminus\{y_1\}}}(x,M)\cap B_{j_{\mathbb{R}^n\setminus\{y_2\}}}(x,M)$. This is clear, because the j-distance between a and b depends only on the closest boundary point of the end points a and b. Similarly for $\partial G=\{y_1,y_2,\ldots,y_m\}$ we have

$$B_{j_G}(x,M)=\bigcap_{i=1}^{m}B_{j_{\mathbb{R}^n\setminus\{y_i\}}}(x,M).$$

This gives an idea to prove Theorem 1.1, which shows that j-balls are convex in any domain G for small radius M.



Figure 2: The boundaries of j-disks in a domain with 1, 2, 3 and 6 boundary points.

Proof of Theorem 1.1. Let $x \in G$ be arbitrary. We claim that

$$A = B_{j_G}(x, M) = \bigcap_{z \in \partial G} B_{j_{\mathbb{R}^n \setminus \{z\}}}(x, M) = B. \tag{4.1}$$

Let $y \in B$. We can choose $z' \in \partial G$ with

$$j_{\mathbb{R}^n \setminus \{z'\}}(x, y) = \min_{z \in \partial G} j_{\mathbb{R}^n \setminus \{z\}}(x, y).$$

Because $z' \in \partial G$ we have $j_G(x, y) \le j_{\mathbb{R}^n \setminus \{z'\}}(x, y)$ and therefore $y \in A$.

On the other hand, let $y \in A$. By definition there is a point $z' \in \partial G$ with $\min\{|x - z'|, |y - z'|\} = \min_{z \in \partial G}\{|x - z|, |y - z|\}$. Now $j_{\mathbb{R}^n \setminus \{z'\}}(x, y) \le j_G(x, y)$ and $y \in B$.

By Theorem 3.1 each $B_{j_{\mathbb{R}^n \setminus \{z\}}}(x, M)$ is convex for $0 < M \le \log 2$ and (4.1) $B_{j_G}(x, M)$ is an intersection of convex domains and therefore it is convex.

If $M \in (0, \log 2]$, then $B_{j_G}(x, M)$ is convex and therefore also starlike with respect to x. If $M \in (\log 2, \log(1 + \sqrt{2})]$, then

$$B_j(x, M) = B \setminus \left(\bigcup_{z \in \partial G} A_z \right),$$

where $B = B^n\big(x, (e^M - 1)d(x)\big)$ and $A_z = B^n(c_z z, r_z)$ for $c_z = |z|/(e^M(2 - e^M))$ and $r_z = |z||1 - e^{-M}|/|e^M - 2|$. Let us assume that $B_j(x, M)$ is not strictly starlike with respect to x. Now there exists $a, b \in B$ such that $b \in (x, a)$, $a \in B_j(x, M)$ and $b \notin B_j(x, M)$. Now $b \in B^n(c_z z, r_z)$ for some $z \in \partial G$. By the proof of Theorem 3.4 $a \in B^n(c_z z, r_z)$, which is a contradiction. $\square$

Corollary 4.2. *For a domain $G \subsetneq \mathbb{R}^n$ and $x \in G$ the j-balls $B_j(x, M)$ are simply connected if $M \in \big(0, \log(1 + \sqrt{2})\big]$.*

Proof. By Theorem 1.1 $B_{j_G}(x, M)$ is starlike with respect to x and therefore also simply connected. $\square$

Corollary 4.3. *For a domain $G \subsetneq \mathbb{R}^n$ and $x \in G$ the j-balls $B_j(x, M)$ are strictly convex if $M \in (0, \log 2)$.*

Proof. By the proof of Theorem 1.1 and Theorem 3.1

$$B_j(x, M) = \bigcap_{z \in \partial G} (B_{z,1} \cap B_{z,2}),$$

where $B_{z,i}$ is a Euclidean ball and $x \in B_{z,i}$. Therefore $B_j(x, M)$ is strictly convex. $\square$

Bounds of Theorem 1.1 are sharp as $G = \mathbb{R}^n \setminus \{0\}$ shows. Also the bound $\log(1+\sqrt{2})$ of Corollary 4.2 is sharp. This can be seen by choosing $G = \mathbb{R}^2 \setminus \{0, z\}$ for a certain z and considering $B_j(e_1, M)$ for $M > \log(1+\sqrt{2})$. By the proof of Theorem 3.1 we know that

$$B_j(e_1, M) = B^2(e_1, r_1) \setminus B^2(c, r_2)$$

for $r_1 = e^M - 1$, $c = e_1/\big(e^M(2-e^M)\big)$ and $r_2 = (e^M-1)/\big(e^M(e^M-2)\big)$. Let l be the tangent line of $B^2(c, r_2)$ that goes through e_1. Denote $\{y\} = S^1(c, r_2) \cap l$. Choose z to be the reflection of 0 in the line l. By a simple computation we have

$$|y - e_1| = \frac{e^M - 1}{\sqrt{e^M(e^M - 2)}} < r_1.$$

Let us denote by c' the reflection of c in the line l. Now $B_{j_{\mathbb{R}^2 \setminus \{0,z\}}}(e_1, M) = B^2(e_1, r_1) \setminus \big(B^2(c, r_2) \cup B^2(c', r_2)\big)$ and therefore $B_j(e_1, M)$ is disconnected for $M > \log(1+\sqrt{2})$.

Similar counterexamples can be constructed for $n > 2$. Let us assume $n \geq 2$ and $M > \log(1+\sqrt{2})$. Now we choose

$$G = \mathbb{R}^n \setminus \big(S^{n-1}(z, |z|) \setminus B^n(e_1, 1)\big),$$

where $z \in S^{n-1}(e_1, e^M - 1)$ and the line $[z, e_1]$ is a tangent of $S^{n-1}(c, r)$ for $c = e_1/\big(e^M(2 - e^M)\big)$ and $r = |1 - e^M|/|e^M(2 - e^M)|$. Let $y \in [z, e_1] \cap S^{n-1}(e_1, e^M - 1)$. Now $j_G(e_1, y) = M$ and $j_G\big(e_1, \frac{1}{2}(z + y)\big) < M$. Therefore $B_j(e_1, M)$ is disconnected.

Remark 4.4. The idea of the proof of Theorem 1.1 cannot be used for the quasihyperbolic metric. We always have

$$D_G(x, M) \subset \bigcap_{z \in \partial G} D_{\mathbb{R}^n \setminus \{z\}}(x, M)$$

but inclusion in the other direction is not always true. For example $G = \mathbb{R}^n \setminus \{0, e_1\}$, $x = e_1/4$ and $M = 1$ gives an counterexample. Now $y = e_1(1 - 1/e)$ is on the boundary $\partial D_G(x, M)$ because

$$k_G(x, y) = k_{\mathbb{R}^n\setminus\{0\}}(x, e_1/2) + k_{\mathbb{R}^n\setminus\{e_1\}}(e_1/2, y) = \log 2 + \log(e/2) = 1.$$

On the other hand, $z = e_1\big(1-3/(4e)\big)$ belongs to the boundary $\partial D_{\mathbb{R}^n\setminus\{e_1\}}(x, M)$. Now $0.632 \approx |y| < |z| \approx 0.724$ and therefore $D_{\mathbb{R}^n\setminus\{0\}}(x, M) \cap D_{\mathbb{R}^n\setminus\{e_1\}}(x, M) \not\subset D_G(x, M)$.

The next theorem states convexity of j-balls in convex domains.

Theorem 4.5. *Let $M > 0$, $G \subsetneq \mathbb{R}^n$ be a convex domain and $x \in G$. Then j-balls $B_j(x, M)$ are convex.*

Proof. By Theorem 1.1 we need to consider only the case $M > \log 2$. Let us divide G into two parts $D_1 = \{z \in G \colon d(z) \geq d(x)\}$ and $D_2 = G \setminus D_1$. We will first show that convexity of G implies convexity of D_1. Let us assume that D_1 is not convex. There exists $a, b \in D_1$ such that $c = (a + b)/2 \notin D_1$ and $d(a) = d(x) = d(b)$. Now $B^n\big(a, d(x)\big)$ and $B^n\big(b, d(x)\big)$ does not contain any points of ∂G, but $B^n(c, r)$ for some $r < d(x)$ contains at least one point of ∂G. Therefore G is not convex, which is a contradiction.

Let us consider $B_j(x, M) \cap D_1$. By definition of the j-metric we have for $y \in \partial B_j(x, M) \cap D_1$

$$|x - y| = d(x)\big(e^M - 1\big)$$

and therefore $\partial B_j(x, M) \cap D_1$ is a subset of $S^{n-1}(x, r)$, where $r = d(x)\big(e^M - 1\big)$. By convexity of D_1 the domain $B_j(x, M) \cap D_1$ is convex.

Let us then show that each chord with end points in $B_j(x, M) \cap D_2$ is contained in $B_j(x, M)$. By definition for $y \in \partial B_j(x, M) \cap D_2$ we have

$$d(y) = \frac{|x - y|}{e^M - 1}. \tag{4.6}$$

Let us assume $y_1, y_2 \in B_j(x, M) \cap D_2$ and $z = (y_1 + y_2)/2 \notin B_j(x, M)$. If $z \in D_1$, then $z \in B_j(x, M)$ because $B_j(x, M) \subset B^n(x, r)$. Therefore we may assume $z \in D_2 \setminus B_j(x, M)$. By (4.6) we have $d(y_i) > |x - y_i|/(e^M - 1)$ for $i \in \{1, 2\}$ and $d(z) < |x - z|/(e^M - 1)$. Since $M > \log 2$ we have $c = 1/(e^M - 1) < 1$. Now the boundary ∂G is outside $B^n(y_1, c|x - y_1|) \cup B^n(y_2, c|x - y_2|)$ and has a point in $B^n(z, c|x - z|)$, see Figure 3.

We will show that for $c < 1$ the domain G is not convex. Let us denote by l a line that is a tangent to balls $B^n(y_1, c|x - y_1|)$ and $B^n(y_2, c|x - y_2|)$. Because $d(y_i, l) = c|x - y_i|$ for $i \in \{0, 1\}$ we have

$$d(z, l) = \frac{c|x - y_1| + c|x - y_2|}{2}. \tag{4.7}$$



Figure 3: Line l, Euclidean balls $B_1 = B^n(y_1, c|x-y_1|)$ and $B_2 = B^n(y_2, c|x-y_2|)$ and points y_1, y_2 and z.

By the triangle inequality

$$|x-z| = \left|\frac{x-y_1}{2} + \frac{x-y_2}{2}\right| \leq \frac{|x-y_1|}{2} + \frac{|x-y_2|}{2}$$

and by (4.7)

$$d(z,l) = \frac{c}{2}(|x-y_1| + |x-y_2|) \geq c|x-z|.$$

Now the domain G is not convex, which is a contradiction, and each chord with end points in $B_j(x,M) \cap D_2$ is contained in $B_j(x,M)$.

Since each chord with end points in $B_j(x,M) \cap D_2$ is contained in $B_j(x,M)$, $B_j(x,M) \cap D_2 \subset B^n(x,r)$, D_1 is convex and $\partial B_j(x,M) \cap D_1 \subset S^{n-1}(x,r)$ the j-ball $B_j(x,M)$ is convex. $\square$

Theorem 4.8. *Let $M > 0$ and $G \subsetneq \mathbb{R}^n$ be a starlike domain with respect to $x \in G$. Then the j-balls $B_j(x,M)$ are starlike with respect to x.*

Proof. By Theorem 1.1 we need to consider $M > \log(\sqrt{2}+1)$ which is equivalent to $e^M - 1 > \sqrt{2}$. Let us divide G into two parts $D_1 = \{z \in G \colon d(z) \geq d(x)\}$ and $D_2 = G \setminus D_1$.

Similarly as in the proof of Theorem 4.5 the boundary $\partial B_j(x,M) \cap D_1$ is a subset of a sphere $S^{n-1}(x,r)$ and $B_j(x,M) \subset S^{n-1}(x,r)$. Therefore it is sufficient to show that for each $y \in B_j(x,M) \cap D_2$ the line segment $[x,y]$ is in $B_j(x,M)$.

We will show that all chords $[x,y]$ for $y \in B_j(x,M) \cap D_2$ are contained in $B_j(x,M)$. Let us assume, on the contrary, that there exists points $y_1, y_2 \in \big(\partial B_j(x,M)\big) \cap D_2$ with $y_1 \in (x,y_2)$ and $z = (y_1+y_2)/2 \notin \overline{B_j}(x,M)$. Let us first assume $z \in D_1$. Now $j_G(x,z) > j_G(x,y_2)$ is equivalent to $|x-z|/d(x) >$

$|x-y_2|/d(y_2)$. By the selection of y_1 and y_2 we have $|x-z|<|x-y_2|$ and $d(x)>d(y_2)$ implying $|x-z|/d(x)<|x-y_2|/d(y_2)$, which is a contradiction.

Let us then assume $z \in D_2$. Now

$$\frac{|x-y_1|}{d(y_1)} = \frac{|x-y_2|}{d(y_2)} = e^M - 1 < \frac{|x-z|}{d(z)}$$



Figure 4: Selection of points y_1 and y_2. Gray circles are $B^n\big(y_1,d(y_1)\big)$, $B^n\big(z,d(z)\big)$ and $B^n\big(y_2,d(y_2)\big)$.

and therefore the boundary ∂G does not intersect $B^n\big(y_1,d(y_1)\big)$ or $B^n\big(y_2,d(y_2)\big)$ and contains a point in $B^n\big(z,d(z)\big)$, see Figure 4. This means that G is not starlike with respect to x, which is a contradiction. $\square$

Remark 4.9. (1) Let us consider the domain $G = B^n(0,1) \cup B^n(e_1,1/4) \cup B^n(2e_1,1)$ and show that the j-ball $B = B_j(0,\log 3)$ is connected but the j-sphere $S = \{z \in G \colon j_G(0,z) = \log 3\}$ is disconnected. We have

$$j_G(0,e_1) = \log\left(1+\frac{1}{1/4}\right) = \log 5$$

and therefore all points $x \in G$ with $x_1 = 1$ are neither in B nor on the boundary ∂B. We also have $B, \partial B \subset B^n(0,1) \cup B^n(2e_1,1)$. For all $y \in B^n(2e_1,1) \setminus \{u \in G \colon \angle 0\,2e_1\,u < \operatorname{atan}(1/4)\}$ we have

$$j_G(0,y) = \log\left(1+\frac{|y|}{1-|2-y|}\right) \geq \log(1+2) = \log 3,$$

because $|y| + 2|2-y| \geq 2$. For all $y \in B^n(2e_1,1) \cap \{u \in G \colon \angle 0\,2e_1\,u < \operatorname{atan}(1/4)\}$ we have

$$j_G(0,y) = \log\left(1+\frac{|y|}{d(y)}\right) \geq \log\left(1+\frac{|y_1|}{d(y_1)}\right) \geq \log(1+2) = \log 3$$

and therefore $B \subset B^n(0,1)$ and it is connected.

Let us now consider S and denote $z \in S$. If $z \in B^n(2e_1, 1)$, then $z = 2e_1$. If $z \in B^n(0,1)$, then $z \in \partial B$. Now $S = \partial B \cup \{2e_1\}$ and it is disconnected. In particular, we see that

$$\overline{\{z \in G \colon j_G(0,z) < \log 3\}} \neq \{z \in G \colon j_G(0,z) \leq \log 3\}.$$

(2) We have seen that in convex domains the j-balls are convex and in starlike domains the j-balls are starlike. However in simply connected domains the j-balls need not be simply connected. Let us consider $G = B^n(0,1) \cup B^n(e_1, h) \cup B^n(2e_1, 1)$ for $h \in (0,1)$. Clearly G is simply connected. Let us consider $B = B_j(0, \log 4)$. We have

$$j_G(0, 2e_1) = \log\left(1 + \frac{2}{1}\right) = \log 3$$

and therefore $2e_1 \in B$. Let $x = (x_1, \ldots, x_n) \in G$ with $x_1 = 1$. Now

$$j_G(0,x) \geq j_G(0,e_1) = \log\left(1 + \frac{1}{h}\right)$$

and $x \notin B$ for $h < 1/3$. For $h = 1/4$ the j-ball B is not even connected. Instead of the radius $\log 4$ we could choose any $r > \log 3$.

Questions 4.10. We pose some open questions concerning the quasihyperbolic metric and quasihyperbolic balls.

(1) Is it true that for any domain $G \subsetneq \mathbb{R}^n$ and $x \in G$ the quasihyperbolic ball $D_G(x, M)$ is strictly convex if $M \in (0,1]$?

(2) Is it true that for any domain $G \subsetneq \mathbb{R}^n$ and $x \in G$ the quasihyperbolic ball $D_G(x, M)$ is strictly starlike with respect to x if $M \in (0, \kappa]$ for $\kappa \approx 2.83297$?

(3) Are the quasihyperbolic geodesics unique in every simply connected domain $G \subsetneq \mathbb{R}^2$?

For the case $\mathbb{R}^n \setminus \{0\}$ see Remarks 3.2 and 3.6.

Acknowledgements. This paper is part of the author's PhD thesis, currently written under the supervision of Prof. M. Vuorinen and supported by the Academy of Finland project 8107317.

References

[GO] F.W. Gehring, B.G. Osgood: *Uniform domains and the quasihyperbolic metric.* J. Anal. Math. 36 (1979), 50-74.

[GP] F.W. Gehring, B.P. Palka: *Quasiconformally homogeneous domains.* J. Anal. Math. 30 (1976), 172-199.

[H] P. Hästö: *Gromov hyperbolicity of the j_G and $\tilde{j}_G$ metrics.* Proc. Amer. Math. Soc.134 (2006), no. 4, 1137–1142.

[HIL] P. Hästö, Z. Ibragimov, H. Lindén: *Isometries of relative metrics.* Comput. Methods Funct. Theory 6 (2006), no. 1, 15–28.

[HL] P. Hästö, H. Lindén: *Isometries of the half-apollonian metric.* Complex Var. Theory Appl. 49 (2004), 405-415.

[K] R. Klén: *Local Convexity Properties of Quasihyperbolic Balls in Punctured Space.* In preparation.

[L] H. Lindén: *Quasihyperbolic Geodesics and Uniformity in Elementary Domains.* Dissertation, University of Helsinki, 2005, Ann. Acad. Sci. Fenn. Math. Diss. 146 (2005).

[S] P. Seittenranta: *Möbius-invariant metrics.* Math. Proc. Camb. Phil. Soc. 125 (1999), 511-533.

[V] J. Väisälä: *Quasihyperbolic geometry of domains in Hilbert spaces.* Ann. Acad. Sci. Fenn. Math. 32 (2007), no. 2, 559–578.

[Vu1] M. Vuorinen: *Capacity densities and angular limits of quasiregular mappings.* Trans. Amer. Math. Soc. 263 (1981), 2, 343-354.

[Vu2] M. Vuorinen: *Conformal invariants and quasiregular mappings.* J. Anal. Math. 45 (1985), 69-115.

[Vu3] M. Vuorinen: *Conformal Geometry and Quasiregular Mappings.* Lecture Notes in Math. Vol. 1319, Springer-Verlag, 1988.

[Vu4] M. Vuorinen: *Metrics and quasiregular mappings.* Proc. Int. Workshop on Quasiconformal Mappings and their Applications, IIT Madras, Dec 27, 2005 - Jan 1, 2006, ed. by S. Ponnusamy, T. Sugawa and M. Vuorinen, *Quasiconformal Mappings and their Applications*, Narosa Publishing House, New Delhi, India, 291–325, 2007.

Department of Mathematics
University of Turku
FIN-20014
FINLAND
e-mail: riku.klen@utu.fi